

THE ALTERNATIVE SCHOOL GROUP LTD

Allergy Policy



Author:	HB
Date:	1st September 2024
To be reviewed:	September 2025
Reviewed:	Sept 24;
Version:	2

Introduction

An allergy is a reaction of the body's immune system to substances that are usually harmless. The reaction can cause minor symptoms such as itching, sneezing or rashes but sometimes causes a much more serious reaction called anaphylaxis.

Anaphylaxis is a serious, life-threatening allergic reaction. It is at the extreme end of the allergic spectrum. The whole body is affected often within minutes of exposure to the allergen, but sometimes it can be hours later. Causes can include foods, insect stings, and drugs.

Most healthcare professionals consider an allergic reaction to be anaphylaxis when it involves difficulty breathing or affects the heart rhythm or blood pressure. Anaphylaxis symptoms are often referred to as the ABC symptoms (Airway, Breathing, Circulation).

It is possible to be allergic to anything which contains a protein, however most people will react to a fairly small group of potent allergens.

Common UK Allergens include (but are not limited to):-

Peanuts, Tree Nuts, Sesame, Milk, Egg, Fish, Latex, Insect venom, Pollen and Animal Dander.

This policy sets out how The Alternative School will support pupils and staff with allergies, to ensure they are safe and are not disadvantaged in any way whilst taking part in school life.

1. Role and responsibilities

Parent Responsibilities

- On entry to the school, it is the parent's responsibility to inform the school of any allergies. This information should include all previous serious allergic reactions, history of anaphylaxis and details of all prescribed medication. This can be shared directly with their teacher and on the TAS parent / carer induction pack.
- Parents are to supply a copy of their child's Allergy Action Plan ([BSACI plans](#) preferred) to school. If they do not currently have an Allergy Action Plan this should be developed as soon as possible in collaboration with a healthcare professional e.g. School nurse/GP/allergy specialist.
- Parents are responsible for ensuring any required medication is supplied, in date and replaced as necessary.
- Parents are requested to keep the school up to date with any changes in allergy management. The Allergy Action Plan will be kept updated accordingly.

Staff Responsibilities

- All staff will complete anaphylaxis training through the mandatory outdoor first aid training that is undertaken every three years. The school can request additional staff training for new staff on anaphylaxis training through Sean Whittle, the training manager for outdoor first aid.
- Staff must be aware of the pupils in their care (regular or cover classes) who have known allergies as an allergic reaction could occur at any time and not just at mealtimes. Any food-related activities must be supervised with due caution.
- Staff leading school trips will ensure they carry all relevant emergency supplies. Trip leaders will check that all pupils with medical conditions, including allergies, carry their medication. Pupils unable to produce their required medication will not be able to attend the excursion.
- School Nurse/SENCO/First Aider (*delete or substitute as appropriate*) will ensure that the up-to-date Allergy Action Plan is kept with the pupil's medication.
- It is the parent's responsibility to ensure all medication is in date however the Headteacher will check medication kept at school on a termly basis and send a reminder to parents if medication is approaching expiry.
- The Headteacher keeps a register of pupils who have been prescribed an adrenaline auto-injector (AAI) and a record of use of any AAI(s) and emergency treatment given.

Responsibilities of the person with the allergy

- TAS is aware that some staff might also have allergies that could have serious health consequences for them and will support them in managing their allergy in school and not discriminate against them due to their allergy.
- People are encouraged to have a good awareness of their symptoms and to let an adult know as soon as they suspect they are having an allergic reaction.
- Pupils who are trained and confident to administer their own AAIs will be encouraged to take responsibility for carrying them on their person at all times.

2. Allergy action plans

Allergy action plans are designed to function as individual healthcare plans for children with food allergies, providing medical and parental consent for schools to administer medicines in the event of an allergic reaction, including consent to administer a spare adrenaline auto-injector.

School recommends using the British Society of Allergy and Clinical Immunology ([BSACI Allergy Action Plans](#)) to ensure continuity. This is a national plan that has been agreed by the BSACI, Anaphylaxis UK and Allergy UK.

It is the parent/carer's responsibility to complete the allergy action plan with help from a healthcare professional (e.g. GP/School Nurse/Allergy Specialist) and provide this to the school.

An allergy action plan will be discussed with any staff members who have an allergy. It will be their choice whether they complete an allergy action plan but the school advises that staff do complete one so the school is able to support in the event of an allergic reaction.

4. Emergency Treatment and Management of Anaphylaxis

What to look for:

Symptoms usually come on quickly, within minutes of exposure to the allergen.

Mild to moderate allergic reaction symptoms may include:

- a red raised rash (known as hives or urticaria) anywhere on the body
- a tingling or itchy feeling in the mouth
- swelling of lips, face or eyes
- stomach pain or vomiting.

More serious symptoms are often referred to as the ABC symptoms and can include:

- **AIRWAY** - swelling in the throat, tongue or upper airways (tightening of the throat, hoarse voice, difficulty swallowing).
- **BREATHING** - sudden onset wheezing, breathing difficulty, noisy breathing.
- **CIRCULATION** - dizziness, feeling faint, sudden sleepiness, tiredness, confusion, pale clammy skin, loss of consciousness.

The term for this more serious reaction is anaphylaxis. In extreme cases there could be a dramatic fall in blood pressure. The person may become weak and floppy and may have a sense of something terrible happening. This may lead to collapse and unconsciousness and, on rare occasions, can be fatal.

If the pupil has been exposed to something they are known to be allergic to, then it is more likely to be an anaphylactic reaction.

Anaphylaxis can develop very rapidly, so a treatment is needed that works rapidly. **Adrenaline** is the mainstay of treatment, and it starts to work within seconds.

What does adrenaline do?

- It opens up the airways
- It stops swelling
- It raises the blood pressure

As soon as anaphylaxis is suspected, adrenaline must be administered without delay. Action:

- Keep the child where they are, call for help and do not leave them unattended.
- **LIE CHILD FLAT WITH LEGS RAISED** – they can be propped up if struggling to breathe but this should be for as short a time as possible.
- **USE ADRENALINE AUTO-INJECTOR WITHOUT DELAY** and note the time given. AAls should be given into the muscle in the outer thigh. Specific instructions vary by brand – always follow the instructions on the device.
- **CALL 999** and state **ANAPHYLAXIS (ana-fil-axis)**.
- If no improvement after 5 minutes, administer second AAI.
- If no signs of life commence CPR.
- Call parent/carer as soon as possible.

Whilst you are waiting for the ambulance, keep the child where they are. Do not stand them up, or sit them in a chair, even if they are feeling better. This could lower their blood pressure drastically, causing their heart to stop.

All pupils must go to hospital for observation after anaphylaxis even if they appear to have recovered as a reaction can reoccur after treatment.

5. Supply, storage and care of medication

Depending on their level of understanding and competence, pupils will be encouraged to take responsibility for and to carry their own **two** AAls on them at all times (in a suitable bag/container).

For younger children or those not ready to take responsibility for their own medication, there should be an anaphylaxis kit which is kept safely, not locked away and **accessible to all staff**.

Medication should be stored in a suitable container and clearly labelled with the pupil's name. The pupil's medication storage container should contain:

- Two AAls i.e. EpiPen® or Jext® or Emerade®
- An up-to-date allergy action plan
- Antihistamine as tablets or syrup (if included on allergy action plan)
- Spoon if required
- Asthma inhaler (if included on allergy action plan).

It is the responsibility of the child's parents to ensure that the anaphylaxis kit is up-to-date and clearly labelled, however the Head Teacher will check medication kept at school on a termly basis and send a reminder to parents if medication is approaching expiry.

Parents can subscribe to expiry alerts for the relevant AAls their child is prescribed, to make sure they can get replacement devices in good time.

Older children and medication

Older children and teenagers should, whenever possible, assume responsibility for their emergency kit under the guidance of their parents. However, symptoms of anaphylaxis can come on **very suddenly**, so school staff need to be prepared to administer medication if the young person cannot.

Storage

AAls should be stored at room temperature, protected from direct sunlight and temperature extremes.

Disposal

AAls are single use only and must be disposed of as sharps. Used AAls can be given to ambulance paramedics on arrival or can be disposed of in a pre-ordered sharps bin. Sharps bins to be obtained from and disposed of at a chemist.

If anaphylaxis is suspected **in an undiagnosed individual** call the emergency services and state you suspect ANAPHYLAXIS. Follow advice from them as to whether administration of the spare AAI is appropriate.

6. Staff Training

The named staff members responsible for co-ordinating staff anaphylaxis training and the upkeep of the school's anaphylaxis policy are:-

Ann Flynn

Kirsty Swierkowski

All staff will complete face to face, 16 hour first aid training at the earliest opportunity they begin work at TAS and training is also available on an ad-hoc basis for any new members of staff.

Training includes:

- Knowing the common allergens and triggers of allergy
- Spotting the signs and symptoms of an allergic reaction and anaphylaxis. Early recognition of symptoms is key, including knowing when to call for emergency services

- Administering emergency treatment (including AAls) in the event of anaphylaxis – knowing how and when to administer the medication/device
- Measures to reduce the risk of a child having an allergic reaction e.g. allergen avoidance, knowing who is responsible for what
- Managing allergy action plans and ensuring these are up to date
- A practical session using trainer devices (these can be obtained from the manufacturers' websites: www.epipen.co.uk and www.jext.co.uk and www.emerade-bausch.co.uk)

7. Inclusion and safeguarding

The Alternative School is committed to ensuring that all children with medical conditions, including allergies, in terms of both physical and mental health, are properly supported in school so that they can play a full and active role in school life, remain healthy and achieve their academic potential.

8. Catering

All food businesses (including school caterers) must follow the Food Information Regulations 2014 which states that allergen information relating to the 'Top 14' allergens must be available for all food products.

The Headteacher will inform the Catering Manager of pupils and staff with food allergies.

- Bottles, other drinks and lunch boxes provided by parents for pupils with food allergies should be clearly labelled with the name of the child for whom they are intended.
- The pupil should be taught to also check with catering staff, before selecting their lunch choice.
- Where food is provided by the school, staff should be educated about how to read labels for food allergens and instructed about measures to prevent cross contamination during the handling, preparation and serving of food. Examples include: preparing food for children with food allergies first; careful cleaning (using warm soapy water) of food preparation areas and utensils. For further information, parents/carers are encouraged to liaise with the Catering Manager.
- Food should not be given to primary school age food-allergic children without parental engagement and permission (e.g. birthday parties, food treats).
- Use of food in crafts, cooking classes, science experiments and special events (e.g. fetes, assemblies, cultural events) needs to be considered and may need to be restricted/risk assessed depending on the allergies of particular children and their age.

9. School trips

Staff leading school trips will ensure they carry all relevant emergency supplies. Trip leaders will check that all pupils with medical conditions, including allergies, carry their medication. Pupils unable to produce their required medication will not be able to attend the excursion.

All the activities on the school trip will be risk assessed to see if they pose a threat to allergic pupils and alternative activities planned to ensure inclusion.

Overnight school trips should be possible with careful planning and a meeting for parents with the lead member of staff planning the trip should be arranged. Staff at the venue for an overnight school trip should be briefed early on that an allergic child is attending and will need appropriate food (if provided by the venue).

Sporting Excursions

Children with allergies should have every opportunity to attend sports trips to other schools.

Most parents are keen that their children should be included in the full life of the school where possible, and the school will need their co-operation with any special arrangements required.

10. Allergy awareness and nut bans

The Alternative School supports the approach advocated by Anaphylaxis UK towards nut bans/nut free schools. They would not necessarily support a blanket ban on any particular allergen in any establishment, including in schools. This is because nuts are only one of many allergens that could affect pupils, and no school could guarantee a truly allergen free environment for a child living with food allergy. They advocate instead for schools to adopt a culture of allergy awareness and education.

A 'whole school awareness of allergies' is a much better approach, as it ensures teachers, pupils and all other staff are aware of what allergies are, the importance of avoiding the pupils' allergens, the signs & symptoms, how to deal with allergic reactions and to ensure policies and procedures are in place to minimise risk.

11. Risk Assessment

The Alternative School will conduct a detailed individual risk assessment for all new joining pupils with allergies and any pupils newly diagnosed, to help identify any gaps in our systems and processes for keeping allergic children safe.

12. Useful Links

Anaphylaxis UK Safer Schools Programme -
<https://www.anaphylaxis.org.uk/education/safer-schools-programme/>

AllergyWise for Schools (including certificate) online training -
<https://www.allergywise.org.uk/p/allergywise-for-schools1>

BSACI Allergy Action Plans - <https://www.bsaci.org/professional-resources/resources/paediatric-allergy-action-plans/>

Spare Pens in Schools - <http://www.sparepensinschools.uk>

Department for Education Supporting pupils at school with medical conditions -
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/803956/supporting-pupils-at-school-with-medical-conditions.pdf

Department of Health Guidance on the use of adrenaline auto-injectors in schools -
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/645476/Adrenaline_auto_injectors_in_schools.pdf

Food allergy quality standards (The National Institute for Health and Care Excellence, March 2016) <https://www.nice.org.uk/guidance/qs118>

Anaphylaxis: assessment and referral after emergency treatment (The National Institute for Health and Care Excellence, 2020)
<https://www.nice.org.uk/guidance/cg134?unlid=22904150420167115834>

This child has the following allergies:

Name: _____

DOB: _____

Photo

Mild/moderate reaction:

- Swollen lips, face or eyes
- Itchy/tingling mouth
- Hives or itchy skin rash
- Abdominal pain or vomiting
- Sudden change in behaviour

Action to take:

- Stay with the child, call for help if necessary
- Locate adrenaline autoinjector(s)
- Give antihistamine: _____
(If vomited, can repeat dose)
- Phone parent/emergency contact

● Watch for signs of ANAPHYLAXIS (life-threatening allergic reaction)

Anaphylaxis may occur without skin symptoms: ALWAYS consider anaphylaxis in someone with known food allergy who has **SUDDEN BREATHING DIFFICULTY**

- | | | |
|---|--|---|
| A AIRWAY | B BREATHING | C CONSCIOUSNESS |
| <ul style="list-style-type: none"> • Persistent cough • Hoarse voice • Difficulty swallowing • Swollen tongue | <ul style="list-style-type: none"> • Difficult or noisy breathing • Wheeze or persistent cough | <ul style="list-style-type: none"> • Persistent dizziness • Pale or floppy • Suddenly sleepy • Collapse/unconscious |

IF ANY ONE (OR MORE) OF THESE SIGNS ABOVE ARE PRESENT:

- 1 Lie child flat with legs raised** (if breathing is difficult, allow child to sit)
 -  ✓
 -  ✓
 -  ✗
- 2 Immediately dial 999** for ambulance and say ANAPHYLAXIS ("ANA-FIL-AX-IS")
- 3 In a school with "spare" back-up adrenaline autoinjectors, ADMINISTER the SPARE AUTOINJECTOR** if available
- 4 Commence CPR** if there are no signs of life
- 5 Stay with child** until ambulance arrives, do **NOT** stand child up
- 6 Phone parent/emergency contact**

*** IF IN DOUBT, GIVE ADRENALINE ***

You can dial 999 from any phone, even if there is no credit left on a mobile. Medical observation in hospital is recommended after anaphylaxis. For more information about managing anaphylaxis in schools and "spare" back-up adrenaline autoinjectors, visit sparepensinschools.uk

Emergency contact details:

1) Name: _____



2) Name: _____



Additional instructions:

If wheezy: DIAL 999 and GIVE ADRENALINE using a "back-up" adrenaline autoinjector if available, then use asthma reliever (blue puffer) via spacer

Parental consent: I hereby authorise school staff to administer the medicines listed on this plan, including a "spare" back-up adrenaline autoinjector (AAI) if available, in accordance with Department of Health Guidance on the use of AAIs in schools

Signed: _____

Print name: _____

Date: _____

For more information about managing anaphylaxis in schools and "spare" back-up adrenaline autoinjectors, visit sparepensinschools.uk

This BSACI Action Plan for Allergic Reactions is for children and young people with mild food allergies, who need to avoid certain allergens. For children at risk of anaphylaxis and who have been prescribed an adrenaline autoinjector device, there are BSACI Action Plans which include instructions for adrenaline autoinjectors. These can be downloaded at bsaci.org

For further information, consult NICE Clinical Guidance CG116 Food allergy in children and young people at guidance.nice.org.uk/CG116

This is a medical document that can only be completed by the child's healthcare professional. It must not be altered without their permission. This document provides medical authorisation for schools to administer a "spare" adrenaline autoinjector in the event of the above-named child having anaphylaxis (as permitted by the Human Medicines (Amendment) Regulations 2017). The healthcare professional named below confirms that there are no medical contra-indications to the above-named child being administered an adrenaline autoinjector by school staff in an emergency. **This plan has been prepared by:**

Sign & print name: _____

Hospital/Clinic: _____



Date: _____